

EPHEDRINE SULFATE- ephedrine sulfate injection, solution

Sandoz Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

EPHEDRINE SULFATE

Injection, USP

(50 mg/mL)

Preservative-Free

Single Dose Vial


Protect from light.

Keep vials in tray until time of use.

DESCRIPTION

Ephedrine Sulfate Injection, USP is a sterile, nonpyrogenic solution containing ephedrine sulfate 50 mg/mL in water for injection. It is administered by subcutaneous, intramuscular or intravenous injection as an adrenergic agent. The solution contains no bacteriostat, antimicrobial agent or added buffer. The pH is 5.3 (4.5 to 7.0). The osmolar concentration of the 5% solution is 0.35 mOsmol/mL (calc.).

Ephedrine Sulfate, USP is a sympathomimetic amine chemically designated α -[1-(methylamino) ethyl] benzenemethanol sulfate (2:1)(salt). It has the following structural formula:



CLINICAL PHARMACOLOGY

Therapeutic doses of ephedrine produce mainly relaxation of smooth muscle and, if norepinephrine stores are intact, cardiac stimulation and increased systolic and usually increased diastolic blood pressure. Its vasopressor effect results largely from increased cardiac output and to a lesser extent from peripheral vasoconstriction. Pressor responses to parenteral ephedrine are slower but more prolonged than those produced by epinephrine. Ephedrine stimulates both alpha and beta receptors and its peripheral actions are due partly to norepinephrine release and partly to direct effect on receptors. Ephedrine may deplete norepinephrine stores in sympathetic nerve endings, so that tachyphylaxis to cardiac and pressor effects of the drug may develop. Central nervous system effects are similar to those of amphetamine drugs but less pronounced. The central effects of ephedrine are overshadowed to a large extent by its peripheral actions.

Glycogenolysis in the liver is increased by ephedrine but not as much as by epinephrine; usual doses of ephedrine are unlikely to produce hyperglycemia. Ephedrine increases oxygen consumption and metabolic rate as a probable result of central stimulation.

Ephedrine is rapidly and completely absorbed following parenteral injection. Pressor and cardiac responses to ephedrine persist for one hour following intramuscular or subcutaneous administration of

25 to 50 mg.

Small amounts of ephedrine are slowly metabolized in the liver; metabolites have been identified as p-hydroxyephedrine, p-hydroxynorephedrine, norephedrine, and conjugates of these compounds. The drug and its metabolites are excreted in the urine, mostly as unchanged ephedrine. Rate of urinary excretion is dependent on urinary pH. Percentage excretion of the drug and its metabolites is increased by acidification of the urine. Elimination half-life of the drug has been reported to be about three hours when the urine is acidified to pH 5 and about six hours when urinary pH is 6.3.

INDICATIONS AND USAGE

Ephedrine Sulfate Injection, USP is indicated primarily to counteract the hypotensive effects of spinal or other types of nontopical conduction anesthesia. It is also useful as a pressor agent in hypotensive states following sympathectomy, or following overdose with ganglionic-blocking agents, antiadrenergic agents, veratrum alkaloids or other drugs used for lowering blood pressure in the treatment of arterial hypertension. The drug is sometimes injected to relieve acute bronchospasm, but it is less effective than epinephrine for this purpose.

CONTRAINDICATIONS

Ephedrine is contraindicated in patients with known hypersensitivity to sympathomimetic amines and in patients with angle closure glaucoma. It should not be used in patients anesthetized with agents such as cyclopropane or halothane as these agents may sensitize the heart to the arrhythmic action of sympathomimetic drugs.

Ephedrine should not ordinarily be used in those cases where vasopressor drugs may be contraindicated, e.g., in thyrotoxicosis, diabetes, in obstetrics when maternal blood pressure is in excess of 130/80 and in hypertension and other cardiovascular disorders.

WARNINGS

Ephedrine may cause hypertension resulting in intracranial hemorrhage. Ephedrine may induce anginal pain in patients with coronary insufficiency or ischemic heart disease. The drug also may induce potentially fatal arrhythmias in patients with organic heart disease or who are receiving drugs that sensitize the myocardium. See CONTRAINDICATIONS.

Initially, parenterally administered ephedrine may produce constriction of renal blood vessels and decreased urine formation.

PRECAUTIONS

Ephedrine Sulfate Injection, USP is subject to oxidation and should be protected against exposure to light.

Do not administer unless solution is clear and seal is intact. Discard unused portion.

Ephedrine should be used cautiously in patients with hyperthyroidism, hypertension, heart disease (including coronary insufficiency, angina pectoris and patients receiving digitalis), cardiac arrhythmias, diabetes or unstable vasomotor system. All vasopressors should be used cautiously in patients taking monoamine oxidase (MAO) inhibitors.

Ephedrine should not be administered concomitantly with other sympathomimetic drugs because of possible additive effects and increased toxicity.

Alpha-adrenergic blocking agents may reduce the vasopressor response to ephedrine by causing vasodilation.

Beta-adrenergic blocking drugs may block the cardiac and bronchodilating effects of ephedrine.

Administration of ephedrine to patients receiving anesthesia with cyclopropane or halogenated hydrocarbons such as halothane which sensitize the myocardium, may induce cardiac arrhythmia. (See CONTRAINDICATIONS). Use of a pressor drug with less cardiac stimulating effects should be considered in patients receiving myocardial sensitizing anesthetics. When encountered, such arrhythmias may respond to administration of a beta-adrenergic blocking drug.

Ephedrine also should be used cautiously with other drugs (e.g., digitalis glycosides) that sensitize the myocardium to the actions of sympathomimetic agents.

Drugs such as reserpine and methyldopa which reduce the amount of norepinephrine in sympathetic nerve endings may reduce the pressor response to ephedrine. Diuretic agents also may decrease vascular response to pressor drugs such as ephedrine.

Ephedrine may antagonize the neuron blockade produced by guanethidine resulting in decreased anti-hypertensive effect and requiring increased dosage of the latter.

Pregnancy Category C

Animal reproduction studies have not been conducted with ephedrine. It is also not known whether ephedrine can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Ephedrine should be given to a pregnant woman only if clearly needed.

Labor and Delivery

Parenteral administration of ephedrine to maintain blood pressure during low or other spinal anesthesia for delivery can cause acceleration of fetal heart rate and should not be used in obstetrics when maternal blood pressure exceeds 130/80. See CONTRAINDICATIONS.

Pediatric Use

The safety and effectiveness of Ephedrine has not been established. Its limited use in pediatric patients has been inadequate to fully define the proper dosage and limitations of use.

ADVERSE REACTIONS

Acute toxic effects are usually extensions of the therapeutic actions of the drug and are most often due to overdosage. Excessive doses may cause a sharp rise in blood pressure sufficient to produce cerebral hemorrhage. Other effects (usually transient) include headache, restlessness, anxiety, tension, tremor, weakness, dizziness, confusion, delirium hallucinations, pallor, respiratory difficulty, palpitation, sweating, nausea or vomiting. Repeated injections may cause contraction of the bladder sphincter and interfere with voluntary urination. The possibility of urinary retention, especially in the elderly male, should be kept in mind.

DRUG ABUSE AND DEPENDENCE

None known with parenteral form.

OVERDOSAGE

Continued injections of ephedrine (after depletion of norepinephrine from the nerve endings with loss of vasopressor effect) may result in hypotension more serious than that existing prior to the use of ephedrine. In the absence of norepinephrine depletion, excessive parenteral dosage produces tachycardia, exaggerated rise in blood pressure, and possible cerebrovascular bleeding, plus central nervous system effects. In the event of adverse blood pressure effects, the drug should be stopped and appropriate corrective measures instituted. See ADVERSE REACTIONS.

DOSAGE AND ADMINISTRATION

Depending on the clinical circumstances, Ephedrine Sulfate Injection may be given subcutaneously, intramuscularly or intravenously.

Usual adult dose: 25 to 50 mg (range 10 to 50 mg) injected subcutaneously or intramuscularly (equivalent to 0.2 to 1.0 mL of 5% solution) is usually adequate to prevent or minimize hypotension secondary to spinal anesthesia. Repeat doses should be governed by blood pressure response or, if used as a bronchodilator, according to the degree of improvement. Absorption (onset of action) by the intramuscular route is more rapid (within 10 to 20 minutes) than by subcutaneous injection. The intravenous route may be used if an immediate effect is desired.

When used during labor, administer only sufficient dosage to maintain blood pressure at or below 130/80.

In acute attacks of asthma, the smallest effective dose should be used (usually 0.25 to 0.5 mL) or as otherwise determined by the patient's response.

Usual pediatric dose: 750 micrograms per kg of body weight or 25 mg/M² of body surface injected intravenously or subcutaneously, four times daily or as otherwise determined by the patient's response.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. See PRECAUTIONS.

HOW SUPPLIED

Ephedrine Sulfate Injection, USP (50 mg/mL) is supplied as follows:

SINGLE DOSE VIAL

NDC Number	Volume
66758-008-01	1 mL fill in 2 mL vial
66758-008-02	1 mL fill in 2 mL vial, 25 × 2 mL

Store at controlled room temperature 15° to 30°C (59° to 86°F).

Caution: Federal (USA) law prohibits dispensing without prescription.

For Sandoz Inc. Customer Service, call 1-800-525-8747.

Manufactured for:

SANDOZ

Princeton, NJ 08540

L-029-00

Package Label - Principal Display Panel - 50 mg Carton

SANDOZ


25 × 2 mL Vials NDC 66758-008-02

Ephedrine Sulfate
Injection, USP
50 mg/mL

1 mL fill in a 2 mL Vial
Single Dose Vial

Preservative Free

Rx only



EPHEDRINE SULFATE

ephedrine sulfate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:66758-008
Route of Administration	SUBCUTANEOUS, INTRAMUSCULAR, INTRAVENOUS	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ephedrine sulfate (ephedrine)	ephedrine sulfate	50 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
water	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66758-008-02	25 in 1 CARTON		
1	NDC:66758-008-01	1 mL in 1 VIAL		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		03/26/2004	

Labeler - Sandoz Inc (005387188)

Revised: 5/2010

Sandoz Inc